

Weed & Feed

DOC ID 529307

Note: Product name "LESCO" is part of bag art page 2.

THIS PRODUCT IS A COMBINATION OF HERBICIDE AND TURF FOOD WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Contains LESCO® Poly Plus® Sulfur Coated Urea to provide a uniform growth with extended nitrogen feeding.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*0.560%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid***†0.145%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***†0.143%
OTHER INGREDIENTS: 99.152%
Total100.000%

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid.....0.46%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid0.12%
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid0.12%

†CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p

18-2-3

**MINI FERTILIZER
GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen (N)..... 18.00%
 0.75% Ammoniacal Nitrogen
 17.25% Urea Nitrogen*
Available Phosphate (P₂O₅) 2.00%
Soluble Potash (K₂O) 3.00%
Sulfur (S)Total 2.70%
 2.70% Free Sulfur (S)
Derived From: Polymer Coated Sulfur Coated Urea, Urea, Ammonium Phosphate, Muriate of Potash.
Chlorine (Cl) Max. 2.50%
*4.90% Slowly Available Urea Nitrogen from Polymer Coated Sulfur Coated Urea.
Made in U.S.A. **F1560**

KEEP OUT OF REACH OF CHILDREN CAUTION

For Chemical Spill, Leak, Fire or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using the toilet.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.
User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing /PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor for treatment advice.• Have person sip a glass of water if able to swallow• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow entry to the treated area until dusts have settled.

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial, and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees, and Roughs) and similar turf areas. Not for use on turf being grown for sale or other commercial sod or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near feed or foodstuffs. Do not store near open containers of fertilizer, seed, or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: Pesticides wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

E.P.A. Reg. No. 228-281-10404
E.P.A. Est. No. 82757-OH-001 (M), 82757-FL-001 (S)
First Letter of Lot Code indicates manufacturing site.

#026217 T

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Arichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Bidden, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False Sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleaf marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherd's purse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bulthistle, Western clematiss, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results: 1) Avoid mowing 1 to 2 days before and after application; 2) Apply when grass is moist (after a rain, sprinkling or dew); 3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and 4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

SPREADER SETTINGS: The following settings are approximate for applying contents of this package at a rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 sq ft.

One pass Application:

MAKE	SETTINGS
LESCO Calibration Gauge	#13
SCOTTS® R8A	1½
Cyclone® or Spyker®	3½
LESCO Pendulum	20
Lely®	3½

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(0.72 lbs) pounds of nitrogen per 1,000 square feet (optional Px05 and K20 levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert Fertilizer claimed.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, INC. BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES. LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn. RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Poly Plus is comprised of Polymer Coated Sulfur Coated Urea.

LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. SCOTTS is a registered trademark of the SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC. Lely is a registered trademark of C Van Der Lely N. V.

G:\REGUL\WP\Erievew\Aprvrd-Lbl Word\026217.doc Rev. 7/14/11 VT (100509)

ACCEPTED
FOR REGISTRATION

Dec. 2, 2011

Net Weight 50 lb (22.7 kg)

F1560
Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114


New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

LESCO®

THIS IS REPRESENTATIVE OF THE ARTWORK FOR THE BAG TO WHICH THE ATTACHED LABEL IS AFFIXED. THE SWOOSH DESIGN COLOR VARIES ACCORDING TO TYPE OF PRODUCT:

- YELLOW:** Pre-Emergents
- ORANGE:** Fungicides
- BLUE:** Herbicides, Weed & Feed
- RED:** Insecticides
- BLACK:** Pest Control
- PURPLE:** Non-Selective Herbicides

LESCO®



FREIGHT SHIPPING BAG
Meeting Requirements of
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed by
HOOD PACKAGING CORPORATION
TYLER, TEXAS

18 X 30