



A complete broad-spectrum postemergence herbicide for industrial, turf, ornamental, forestry, roadside, utility rights-of-way, and other listed terrestrial weed control.

(For a complete list of terrestrial use sites, see the Directions for Use section of the label.)

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt . . .	41.0%
OTHER INGREDIENTS (including surfactant):	59.0%
	100.0%

[†]Contains 480 grams of the active ingredient glyphosate, in the form of its isopropylamine salt, per liter or 4 pounds per U.S. gallon, which is equivalent to 356 grams of the acid, glyphosate, per liter or 3 pounds per U.S. gallon (30.4% by weight).

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

Keep out of reach of children

CAUTION

Causes eye irritation. Avoid contact with eyes or clothing.

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
	<ul style="list-style-type: none"> • Have the product container or labeling with you when calling a poison control center or doctor, or going for treatment. • You can also call (314) 694-4000, collect, day or night, for emergency medical treatment information. • This product is identified as Roundup PRO Herbicide, EPA Registration No. 524-475.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment).

If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

See attached labeling for additional Precautions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

See Complete Directions for Use attached to this label for complete Agricultural and Non-Agricultural Use Requirements of the Worker Protection Standard, Directions for Use and Limit of Warranty and Liability in English and Spanish.

STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal. **PESTICIDE STORAGE:** Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. **PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures. **CONTAINER HANDLING AND DISPOSAL:** Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state. Triple rinse or pressure rinse (or equivalent) this container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip. Once properly rinsed, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest collection site, contact your chemical dealer or Monsanto Company at 1-800-ROUNDUP (1-800-768-6387). If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

For product information or assistance using this product call toll-free 1-800-332-3111

THIS IS AN END-USE PRODUCT. MONSANTO COMPANY DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.

EPA Reg. No. 524-475

EPA Est. 524-IA-1

Packed for: MONSANTO COMPANY 800 N. LINDBERGH BLVD. ST. LOUIS, MISSOURI 63167 USA

©2018

101816 ACCEPTED 21136S14-22

DOC ID 557111

VIA NOTIFICATION
LABEL NOT REVIEWED

May 15, 2018

NET 2.5 GAL

MONSANTO
New York State Department

of Environmental Conservation

Division of Materials Management

Pesticide Product Registration

Monsanto Reference ID 2018-NNY166

This page left intentionally blank

2D
CODE



92-075510-26

GROUP 9 HERBICIDE

OPEN
HERE



Complete Directions for Use

A complete broad-spectrum postemergence herbicide for industrial, turf, ornamental, forestry, roadside, utility rights-of-way, and other listed terrestrial weed control.
(For a complete list of terrestrial use sites, see the Directions for Use section of this label.)

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt 41.0%
OTHER INGREDIENTS (including surfactant): 59.0%
100.0%

*Contains 480 grams of the active ingredient glyphosate, in the form of its isopropylamine salt, per liter or 4 pounds per U.S. gallon, which is equivalent to 356 grams of the acid, glyphosate, per liter or 3 pounds per U.S. gallon (30.4% by weight).

Keep out of reach of children

CAUTION See inside for additional precautions.

Causes eye irritation
Avoid contact with eyes or clothing

FIRST AID

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
<ul style="list-style-type: none">• Have the product container or labeling with you when calling a poison control center or doctor, or going for treatment.• You can also call (314) 694-4000, collect, day or night, for emergency medical treatment information.• This product is identified as Roundup PRO Herbicide, EPA Registration No. 524-475.	

Read the entire label before using this product.

Use only according to label directions.

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION, COULD RESULT.

THIS IS AN END-USE PRODUCT. MONSANTO COMPANY DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

EPA Reg. No. 524-475

Packed for: MONSANTO COMPANY
800 N. LINDBERGH BLVD.
ST. LOUIS, MISSOURI, 63167 USA

©2018

020316

	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Perennial Weed Species		
Pepperweed, perennial	4	2
Phragmites*	3–5	1–2

For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts of this product per acre as a broadcast application or a 2-percent solution using a handheld sprayer. In other areas of the U.S., apply 3 to 5 quarts per acre as a broadcast application or, for partial control, apply a 1-percent solution using a handheld sprayer. For enhanced results, make application in late-summer or fall when plants are actively growing and in full bloom. Due to the dense nature of this vegetation (which can prevent good spray coverage) and uneven stages of growth, more than one application of this product might be necessary to achieve control. Visual symptoms of control will be slow to develop.

Quackgrass	2–3	2
Apply this product when most target plants are at least 8 inches in height (3- to 4-leaf stage of development) and actively growing.		
Redvine*	2	2
Reed; common, giant	4–5	2

For enhanced results, make application in late-summer or fall. Control can also be achieved by injecting 8 milliliters of this concentrated product (undiluted) directly into the second or third internode using a handheld injection device.¹

Ryegrass, perennial	2–3	1
----------------------------	-----	---

Apply this product when most target plants have reached the boot to head stage of growth. When applied prior to the boot stage, reduced control can result. In the fall, make application before ryegrass turns brown.

Smartweed, swamp	3–5	2
Spatterdock	4	1

Make application when most target plants are in full bloom. For enhanced results, apply in the summer or fall.

Sowthistle, perennial	2–3	2
Spurge, leafy*	–	2
Startistle, yellow	2–3	2
Sweet potato, wild*	–	2

Make application when most target plants are at or beyond the bloom stage of growth. More than one application will be needed to achieve control.

Thistle, artichoke	2–3	1–2
---------------------------	-----	-----

Make application when target plants are at or beyond the bud stage of growth.

Thistle, Canada	2–3	2
------------------------	-----	---

Make application when target plants are at or beyond the bud stage of growth. Control can also be achieved by stem-injection. Cut 8 to 9 of tallest plants in a clump at bud stage. Push a cavity needle into the stem center and then slowly remove it as you inject 0.6 milliliter of this concentrated product into the stem.¹

Timothy	2–3	2
----------------	-----	---

Make application when most target plants have reached the boot to head stage of development. If application is made prior to the boot stage, reduced control can result. In the fall, make application before plants turn brown.

Torpedgrass*	4–5	2
Trumpet creeper*	2–3	2
Tules, common	–	2

Make application to target plants at or beyond the seedhead stage of development. Visual symptoms will be slow to appear and might not appear for 3 or more weeks after application.

Vaseygrass	3–5	2
Velvetgrass	3–5	2
Wheatgrass, western	2–3	2

Make application when most target plants have reached the boot to head stage of development. Application made prior to the boot stage could result in reduced control. In the fall, make application before plants turn brown.

¹ Partial control only
² When using stem injection, the combined total use of this product must not exceed 10.6 quarts per acre per year. At 6 milliliters of concentrated (undiluted) product per stem, 10.6 quarts will treat approximately 1670 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

11.4 Woody Brush, Trees and Vines

Apply this product to brush and trees that are actively growing after full leaf expansion, unless otherwise directed. Use a higher application rate or spray solution concentration within a given range to control larger brush and trees and/or for application in areas of dense vegetative growth, or for the control of vines that have reached the woody stage of growth.

Enhanced control of woody brush and trees can be obtained when application is made in late-summer or fall after fruit formation; however, in arid areas, enhanced control can be obtained when application is made in the spring to early-summer when brush and trees are at high moisture content and flowering. Poor control can be expected when this product is applied to drought-stressed brush and trees.

Some autumn color on undesirable deciduous species is acceptable when applying this product to brush and trees in the fall, provided no major leaf drop has occurred. Reduced performance of this product could result if application is made following a frost. Symptoms might not appear prior to frost or senescence following a fall application.

For enhanced results, allow 7 or more days after application before mowing, cutting, tilling, burning or removal of woody brush, trees and vines from the application site. Additional applications of this product will be needed to control brush and trees regenerating from underground parts or seed.

TANK MIXTURES: This product may be applied at any rate stated on this label in a tank mixture with the following products to increase the spectrum of control of herbaceous weeds, woody brush, trees and vines. For control of herbaceous weeds, apply the tank-mix product at the lower end of the given application rate or spray solution concentration range. For control of dense stands or hard-to-control woody brush, trees and vines, increase the application rate or spray solution concentration of the tank-mix product towards the higher end of the range. Refer to the individual product labels for approved uses and application rates.

Arsenal; Arsenal Herbicide Applicators Concentrate; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Vastlan Specialty; imazapyr; metsulfuron methyl; triclopyr

Ensure that the proper amount of the Garlon herbicide is thoroughly mixed with water in the spray tank before adding this product.

Cut Stump Application

This product may be used to control re-growth and re-sprouting of woody brush and trees on any site listed on this label.

Cut the woody brush or tree close to the soil surface and immediately apply a 50- to 100-percent (undiluted) solution of this product to the freshly-cut surface using an applicator capable of applying this product to the entire cambium. A delay in application could result in reduced performance. For enhanced results, cut the woody brush or tree during period of active growth and full leaf expansion and apply this product.

For control of the Tree of heaven (*Ailanthus altissima*), cut the tree close to the soil surface and immediately apply a 50-percent solution of this product (16 fluid ounces per quart of solution) and 10 percent Arsenal herbicide (3 to 4 fluid ounces per quart of solution) in water to the freshly-cut surface.

DO NOT MAKE A CUT STUMP APPLICATION WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MIGHT BE GRAFTED TO THE ROOTS OF THE CUT STUMP, AS INJURY COULD OCCUR IN THE ADJACENT TREES. Some sprouts, stems, or trees can share a common root system. Adjacent trees having a similar age, height and spacing could be an indicator of a shared root system. Whether grafted or shared, injury is likely to occur to adjacent stems or trees when this product is applied to one or more trees sharing a common root system.

Woody Brush and Tree Injection and Frill Application

This product may be used to control woody brush and trees listed in this section by injection or frill application on any site listed on this label.

Inject or apply the equivalent of 1 milliliter (0.04 fluid ounce) of this product for every 2 to 3 inches of trunk diameter at breast height (DBH). If injecting this product into the woody brush or tree, use equipment capable of penetrating into the living plant tissue under the bark.

For frill application, apply a 50- to 100-percent (undiluted) solution of this product in water to either a continuous frill around the tree or to cuts evenly spaced around the tree below all branches. As tree diameter increases, enhanced results can be achieved by applying this product to a continuous frill or to more closely spaced cuttings. Avoid application techniques that allow runoff of this product to occur from

frilled or cut areas. In species that freely exude sap, make the frill or cuts at an oblique angle to produce a cupping effect and apply this concentrated product undiluted. For enhanced results, make this application during period of active growth and after full leaf expansion.

Modified High-Volume and Low-Volume Backpack Application

For control and partial control of woody brush, trees and vines listed on this label when using a backpack sprayer or other handheld equipment and a directed low-volume foliar application technique, apply a 5- to 10-percent solution of this product evenly over the plant crown to cover 50 percent of the upper foliage of undesirable vegetation.

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Alder	3–4	1–1.5
Ash ¹	2–5	1–2
Aspen, quaking	2–3	1–1.5
Beardclover (Bearmat) ¹	2–5	1–2
Beech ¹	2–5	1–2
Birch	2	1
Blackberry	3–4	1–1.5
Blackgum	2–5	1–2
Bracken	2–5	1–2
Broom; French, Scotch	2–5	1.5–2
Buckwheat, California ²	2–4	1–2
Cascara ¹	2–5	1–2
Castor bean	2–5	2

Also for control, inject 5 milliliters of this concentrated (undiluted) product per plant directly into the lower portion of the main stem using a handheld injection device.¹

Gatsclaw ²	–	1–1.5
-----------------------	---	-------

For partial control, apply this product when at least 50 percent of the new leaves are fully developed.

Ceanothus ²	2–5	1–2
Chamise ²	2–5	1
Cherry, bitter, black, pin	2–3	1–1.5
Coyote brush	3–4	1.5–2

For control, apply this product when at least 50 percent of the new leaves are fully developed.

Deerweed	2–5	1
Dogwood ²	2–5	1–2
Elderberry	2	1
Elm ²	2–5	1–2
Eucalyptus, blue gum	–	2

For control of eucalyptus re-sprouts, apply this product using a handheld sprayer when re-sprouts are 6 to 12 feet tall. Ensure complete coverage.

Gallberry	2–5	1–2
Gorse ²	2–5	1–2
Hackberry, western	2–5	1–2
Hasardia ²	2–4	1–2
Hawthorn	2–3	1–1.5
Hazel	2	1
Hickory ²	2–5	1–2
Honeysuckle	3–4	1–1.5
Hornbeam, American ²	2–5	1–2
Ivy, poison	3–4	2
Kudzu	4	2
Locust, black ²	2–4	1–2
Madrone (re-sprouts) ¹	–	2
Manzanita ²	2–5	1–2
Maple, red	2–4	1–1.5

For control, apply a 1- to 1.5-percent solution of this product using a handheld sprayer when leaves are fully developed. For partial control, apply 2 to 4 quarts per acre as a broadcast application.

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Maple, sugar	–	1–1.5
For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.		
Maple, vine ²	2–5	1–2
Monkey flower ¹	2–4	1–2
Oak, black, white ¹	2–4	1–2
Oak, northern, pin	2–4	1–1.5
For control, apply this product when at least 50 percent of the new leaves are fully developed.		
Oak, poison	4–5	2
Repeat applications might be required to maintain control. Application in the fall must be made before leaves lose green color.		
Oak, post	3–4	1–1.5
Oak, red	–	1–1.5
For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.		
Oak, scrub ²	2–4	1–1.5
Oak, southern red	2–3	1–1.5
Orange, Osage	2–5	1–2
Peppertree, Brazilian (Florida holly)*	2–5	1–2
Persimmon ²	2–5	1–2
Pine	2–5	1–2
Poplar, yellow ²	2–5	1–2
Redbud, eastern	2–5	1–2
Rose, multiflora	2	1
Make application prior to leaf deterioration by leaf-feeding insects.		
Russian olive ²	2–5	1–2
Sage, black	2–4	1
Sage, white ²	2–4	1–2
Sagebrush, California	2–4	1
Salmonberry	2	1
Saltcedar ²	2–5	1–2
For partial control, apply a 1- to 2-percent solution of this product using a handheld sprayer or 2 to 5 quarts per acre as a broadcast application. For control, apply a 1- to 2-percent solution of this product in a tank-mix with Arsenal herbicide or Arsenal Herbicide Applicators Concentrate using a handheld sprayer. For control using broadcast application, apply 2.25 quarts of this product per acre in a tank-mix with an appropriate rate of Arsenal herbicide or Arsenal Herbicide Applicators Concentrate to plants less than 6 feet tall. To control saltcedar greater than 6 feet tall using broadcast application, apply 4.5 quarts of this product per acre in a tank-mix with a higher rate of Arsenal herbicide or Arsenal Herbicide Applicators Concentrate.		
Sassafras ²	2–5	1–2
Sourwood ²	2–5	1–2
Sumac; laurel, poison, smooth, sugarbush, winged ²	2–4	1–2
Sweetgum	2–3	1–1.5
Swordfern ²	2–5	1–2
Tallowtree, Chinese	–	1
Tan oak (re-sprouts)*	–	2
Thimbleberry	2	1
Tobacco, tree ²	2–4	1–2
Toyon ²	–	2
Trumpet creeper	2–3	1–1.5
Virginia creeper	2–5	1–2
Waxmyrtle, southern ²	2–5	1–2
Willow	3	1
Yerba Santa, California ²	–	2

¹ Partial control

12.0 LIMIT OF WARRANTY AND LIABILITY

Monsanto Company ("Company") warrants that this product conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall use this product only for the purposes of and in accordance with the Complete Directions for Use Label ("Directions") and shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss, injuries or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, crop injury or failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, use and/or application in any manner not explicitly set forth in or inconsistent with the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY TO THE FULLEST EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Certainty, Monsanto and Vine Design, Outrider, Roundup, and Roundup PRO and Design, are registered trademarks of Monsanto Technology LLC.

All other trademarks are the property of their respective owners.

In case of an emergency involving this product, or for medical assistance, call collect, day or night, (314) 694-4000.



©2018
Packed For:
MONSANTO COMPANY
800 N. LINDBERGH BLVD.
ST. LOUIS, MISSOURI 63167 U.S.A.

Roundup PRO[®]

Herbicida

GRUPO 9 HERBICIDA

Instrucciones de uso completas

Un herbicida completo de amplio espectro para aplicación postemergencia, para el control de malezas industrial, ornamental, en céspedes, forestación, lados de carreteras, servidumbres de paso y otros usos terrestres indicados.

(Para una lista completa de usos terrestres, vea la sección Modo de empleo de esta etiqueta.)

INGREDIENTE ACTIVO:

*Glifosato, N-(fosfonometil) glicina, en la forma de su sal de isopropilamina 41.0%
 OTROS INGREDIENTES (incluyendo surfactante): 59.0%
 100.0%

*Contiene 480 gramos por litro o 4 libras por galón norteamericano del ingrediente activo glifosato, en la forma de su sal de isopropilamina, lo cual es equivalente a 356 gramos por litro o 3 libras por galón norteamericano (30.4% por peso) del ácido, glifosato.

Manténgase fuera del alcance de los niños PRECAUCIÓN

Causa irritación a los ojos Evite el contacto con los ojos o la ropa

PRIMEROS AUXILIOS	
SI ENTRA EN CONTACTO CON LOS OJOS	<ul style="list-style-type: none"> Mantenga los ojos abiertos y enjuague lenta y cuidadosamente con agua durante 15 a 20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame a un centro de control de envenenamientos o a un médico para que le indique el tratamiento.
<ul style="list-style-type: none"> Tenga a mano el envase o la etiqueta del producto cuando llame a un centro de control de envenenamientos o a un médico, o cuando vaya a obtener tratamiento. También puede llamar al (314) 694-4000, por cobrar, de día o de noche, para información sobre tratamiento médico de urgencia. Este producto está identificado como Roundup PRO Herbicida, N° de Registro de la EPA 524-475. 	

Lea toda la etiqueta antes de usar este producto.

Use solo según las instrucciones de la etiqueta. EVITE EL CONTACTO DE ESTE HERBICIDA CON EL FOLLAJE, TALLOS VERDES, RAÍCES NO LEÑOSAS EXPUESTAS O FRUTOS EXPUESTOS DE LOS CULTIVOS, PLANTAS Y ÁRBOLES DESEABLES, PORQUE LAS PLANTAS PUEDEN SUFRIR GRAVES DAÑOS O SER DESTRUIDAS.

ESTE ES UN PRODUCTO PARA USARSE TAL Y COMO ESTÁ PREPARADO! MONSANTO COMPANY NO LO HA DISEÑADO NI LO HA REGISTRADO! PARA QUE SEA REFORMULADO. VEA LA ETIQUETA DEL ENVASE INDIVIDUAL PARA CONOCER LAS LIMITACIONES DE REEMPAQUE.

Antes de comprar o usar el producto, lea "LÍMITES EN LA GARANTÍA Y EN LA RESPONSABILIDAD" en la última sección de la etiqueta. Si las condiciones son inaceptables, devuelva el producto inmediatamente, sin abrir el envase.

N° Reg. EPA 524-475
 Envasado para: MONSANTO COMPANY
 800 N. LINDBERGH BLVD.
 ST. LOUIS, MISSOURI, 63167 USA

©2018

020316

No todos los productos recomendados en esta etiqueta han sido registrados para su uso en California. Verifique el estado de registro de cada producto en California antes de utilizarlo.

CONTENIDO

1	1.0	INGREDIENTES	13
2	2.0	TELÉFONOS IMPORTANTES	13
3	3.0	DECLARACIONES PREVENTIVAS	13
	3.1	Riesgos para los seres humanos y los animales domésticos	13
	3.2	Riesgos para el medio ambiente	13
	3.3	Riesgos físicos o químicos	13
4	4.0	ALMACENAMIENTO Y ELIMINACIÓN	13
5	5.0	INFORMACIÓN DEL PRODUCTO	14
6	6.0	MANEJO DE RESISTENCIA DE MALEZAS	14
	6.1	Prácticas de manejo de malezas	14
	6.2	Manejo de biotipos resistentes al glifosato	14
7	7.0	MEZCLAS	14
	7.1	Mezclar con agua	14
	7.2	Mezclas de tanque	14
	7.3	Procedimiento de mezcla en tanque	15
	7.4	Mezclar concentraciones de solución de rocío	15
	7.5	Colorantes y tintes	15
8	8.0	EQUIPOS Y TÉCNICAS PARA LA APLICACIÓN	15
	8.1	Manejo de la dispersión del rocío	15
	8.2	Equipo de aplicación aérea	15
	8.3	Equipo de aplicación terrestre	16
	8.4	Rociadores manuales	16
	8.5	Equipo de aplicación selectiva	16
	8.6	Sistemas por inyección	17
	8.7	Aplicador por goteo controlado (CDA)	17
9	9.0	SITIOS DE USO TERRESTRE	17
10	10.0	INFORMACIÓN ADICIONAL SOBRE GERENCIA DEL LUGAR	17
	10.1	Manejo de bosques, árboles de madera y árboles de Navidad	17
	10.2	Manejo de hábitats de vida silvestre y especies nativas	18
	10.3	Manejo de vivero ornamental y de producción	18
	10.4	Manejo de áreas comerciales, residenciales y recreativas	18
	10.5	Manejo de zonas de pasturas	18
	10.6	Manejo de ferrocarriles	18
	10.7	Manejo de tierras de pastoreo	18
	10.8	Manejo de lados de carreteras	19
	10.9	Manejo de servicios públicos	19
11	11.0	TIPOS DE MALEZAS CONTROLADAS	19
	11.1	Control de malezas, renovación y segado químico en céspedes	19
	11.2	Malezas anuales	20
	11.3	Malezas perennes	20
	11.4	Árboles, enredaderas y matorrales leñosos	22
12	12.0	LÍMITES EN LA GARANTÍA Y EN LA RESPONSABILIDAD	23

1.0 INGREDIENTES

INGREDIENTE ACTIVO:

* Glifosato, N-(fosfonometil) glicina, en la forma de su sal de isopropilamina 41.0%
 OTROS INGREDIENTES (incluyendo surfactante): 59.0%
 100.0%

*Contiene 480 gramos por litro o 4 libras por galón norteamericano del ingrediente activo glifosato, en la forma de su sal de isopropilamina, lo cual es equivalente a 356 gramos por litro o 3 libras por galón norteamericano (30.4% por peso) del ácido, glifosato.

Este producto está protegido por las Patentes de los EE. UU. N° 5,683,958; 5,703,015; 5,063,733; 6,121,199; 6,121,200. No se han otorgado licencias de uso bajo ninguna patente que no sea de los Estados Unidos de América.

2.0 NÚMEROS DE TELÉFONO IMPORTANTES

- PARA INFORMACIÓN SOBRE EL PRODUCTO O AYUDA PARA UTILIZAR ESTE PRODUCTO, LLAME GRATIS AL (800) 332-3111
- EN CASO DE EMERGENCIA RELACIONADA CON ESTE PRODUCTO O PARA AYUDA MÉDICA, LLAME POR COBRAR, DE DÍA O DE NOCHE, AL (314) 694-4000

3.0 DECLARACIONES PREVENTIVAS

3.1 Riesgos para los seres humanos y los animales domésticos

Manténgase fuera del alcance de los niños PRECAUCIÓN

Causa irritación a los ojos Evite el contacto con los ojos o la ropa

PRIMEROS AUXILIOS	
SI ENTRA EN CONTACTO CON LOS OJOS	<ul style="list-style-type: none"> Mantenga los ojos abiertos y enjuague lenta y cuidadosamente con agua durante 15 a 20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame a un centro de control de envenenamientos o a un médico para que le indique el tratamiento.
<ul style="list-style-type: none"> Tenga a mano el envase o la etiqueta del producto cuando llame a un centro de control de envenenamientos o a un médico, o cuando vaya a obtener tratamiento. También puede llamar al (314) 694-4000, por cobrar, de día o de noche, para información sobre tratamiento médico de urgencia. Este producto está identificado como Roundup PRO Herbicida, N° de Registro de la EPA 524-475. 	

ANIMALES DOMÉSTICOS: Se considera que este producto es relativamente no tóxico para perros y otros animales domésticos, sin embargo, la ingestión de este producto o de abundantes cantidades de vegetación rociada recientemente podría causar irritación gastrointestinal temporal (vómitos, diarrea, cólicos, etc.). Si se observan dichos síntomas, dé al animal suficiente cantidad de líquidos para evitar la deshidratación. Llame a un veterinario si los síntomas persisten por más de 24 horas.

Equipo de protección personal (EPP)

Los usuarios y personas que manipulan este producto deben usar: camisas de mangas largas y pantalones largos, además de zapatos y calcetines.

Respete las instrucciones del fabricante para limpiar y mantener los equipos de protección personal (EPP). En caso de que no haya instrucciones, lave el equipo protector con detergente y agua caliente. Mantenga el EPP aparte del resto de la ropa y lávelo por separado.

Recomendaciones de seguridad para el usuario

Los usuarios deben:

- Lavarse las manos antes de comer, beber, masticar chicle, usar tabaco o usar el baño.
- Quitarse la ropa de inmediato si el pesticida traspasa la ropa. Luego deben lavarse muy bien y ponerse ropa limpia.

3.2 Riesgos para el medio ambiente

No aplique directamente al agua, en áreas donde el agua superficial esté presente o en áreas intermareales por debajo del nivel medio de mareas altas. No contamine el agua cuando limpie el equipo o deseche el agua de lavado y enjuague del equipo.

3.3 Riesgos físicos o químicos

Para mezclar, almacenar y aplicar la solución de rocío de este producto, se pueden usar solamente recipientes de acero inoxidable, fibra de vidrio, plástico o recipientes de acero recubiertos internamente con plástico.

NO MEZCLE, ALMACENE O APLIQUE ESTE PRODUCTO O LAS SOLUCIONES DE ROCÍO DE ESTE PRODUCTO EN ENVASES DE ACERO GALVANIZADO O SIN REVESTIMIENTO (EXCEPTO ACERO INOXIDABLE) O EN TANQUES DE ROCÍO. Si se utiliza en estos envases o tanques, este producto o las soluciones de rocío de este producto reaccionan y

producen gas hidrógeno que puede formar una mezcla de gases altamente inflamable. Esta mezcla de gases podría incendiarse o explotar si está en contacto con fuego, chispas, sopletes para soldar, cigarrillos encendidos o cualquier otra fuente de ignición y causar lesiones personales graves.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto de una manera que no sea la indicada en la etiqueta. Este producto solo puede utilizarse de acuerdo con las instrucciones de uso en la etiqueta o según las etiquetas complementarias que se publican por separado. También puede solicitar las etiquetas complementarias a su vendedor minorista autorizado de Monsanto o a su representante de Monsanto Company.

No aplique este producto de manera que entre en contacto con los trabajadores u otras personas, ya sea directamente o por dispersión. Solamente los aplicadores que usan protección podrán estar en el área durante su aplicación. Para verificar requisitos específicos de su tribu o estado, consulte con la agencia responsable de la regulación del uso de pesticidas.

Requisitos para uso agrícola

Utilice este producto solo de acuerdo con la etiqueta y con las Normas de Protección para Trabajadores, 40 CFR Parte 170. Estas Normas contienen los requisitos para la protección de trabajadores agrícolas en granjas, bosques, viveros e invernaderos y para las personas que manipulan pesticidas agrícolas. Contienen los requisitos para capacitar, descontaminar, notificar y ofrecer asistencia de emergencia. También contienen instrucciones específicas y excepciones relativas a las afirmaciones en esta etiqueta sobre los equipos de protección personal (EPP) y los intervalos de acceso restringido. Los requisitos en esta caja se refieren únicamente a las aplicaciones de este producto cubiertas por las Normas de Protección para Trabajadores.

No entre ni permita la entrada de personal a las áreas tratadas durante el intervalo de entrada restringida (REI, por sus siglas en inglés) de 4 horas.

El EPP que se requiere para el acceso anticipado a zonas tratadas de acuerdo con las Normas de Protección para Trabajadores y que incluye el contacto con material tratado, como plantas, tierra o agua es: overoles, zapatos con calcetines y guantes resistentes a sustancias químicas confeccionados con cualquier tipo de material impermeable.

Requisitos para usos no agrícolas

Los requisitos en esta caja se refieren a las aplicaciones de este producto que NO cubren las Normas de Protección para Trabajadores para pesticidas agrícolas (40 CFR, Parte 170). Las Normas se aplican cuando este producto se utiliza para producir plantas agrícolas en granjas, bosques, viveros o invernaderos.

Mantenga a las personas y a las mascotas fuera de las áreas tratadas hasta que la solución de rocío se haya secado.

4.0 ALMACENAMIENTO Y ELIMINACIÓN

El almacenamiento y la eliminación adecuados de los pesticidas son fundamentales para evitar la exposición de las personas y el medio ambiente a consecuencia de pérdidas y derrames del producto, excedentes o desechos o actos de vandalismo. No permita que este producto contamine el agua, ni los alimentos para personas o animales, ni las semillas por medio del almacenamiento o la eliminación.

ALMACENAMIENTO DEL PESTICIDA: Guarde los pesticidas lejos de los alimentos para personas, los alimentos para mascotas, los alimentos para animales, las semillas, los fertilizantes y los materiales de uso veterinario. Mantenga el envase cerrado para evitar los derrames y la contaminación. Vea la etiqueta del envase individual para conocer las condiciones adicionales de almacenamiento, si las hay. **ELIMINACIÓN DEL PESTICIDA:** Para evitar desechos, utilice todo el material contenido en este envase, incluyendo los residuos del enjuague, aplicándolo según las indicaciones de la etiqueta. Si no se pueden evitar los desechos, ofrezca el producto restante a un centro de eliminación de desechos o a un programa de desecho de pesticidas. Estos programas suelen ser manejados por los gobiernos estatales o locales o por la industria. Toda eliminación debe seguir los reglamentos y procedimientos federales, estatales y locales pertinentes.

MANEJO Y ELIMINACIÓN DEL ENVASE: Consulte la etiqueta pegada al envase para conocer las instrucciones de manejo y eliminación del envase, así como los límites de relleno.

Pull at scored cutouts for easier opening.

Pull at scored cutouts for easier opening.

THIS SIDE UP

THIS SIDE UP

2D Code

GROUP 9 HERBICIDE




Herbicide

GROUP 9 HERBICIDE




Herbicide

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product may only be used in accordance with the Directions for Use on this label or in separately published supplemental labeling. Supplemental labeling can be obtained from your Authorized Monsanto Retailer or Monsanto Company Representative.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried.

See attached labeling inside for complete Agricultural and Non-Agricultural Use Requirements of the Worker Protection Standard and Directions for Use in English and Spanish.

For product information or assistance using this product, call toll-free 1-800-332-3111.

Storage and Disposal

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

CONTAINER HANDLING AND DISPOSAL: Nonrefillable container. Do not reuse the container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.

Triple rinse or pressure rinse (or equivalent) the container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest collection site, contact your chemical dealer or Monsanto Company at 1-800-ROUNDUP (1-800-768-6387). If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

LIMIT OF WARRANTY AND LIABILITY

Monsanto Company ("Company") warrants that this product conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall use this product only for the purposes of and in accordance with the Complete Directions for Use label ("Directions") and shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss, injuries or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, crop injury or failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, use and/or application in any manner not explicitly set forth in or inconsistent with the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Monsanto and Vine Design, and Roundup PRO and Design, are registered trademarks of Monsanto Technology LLC.

A complete broad-spectrum postemergence herbicide for industrial, turf, ornamental, forestry, roadside, utility rights-of-way, and other listed terrestrial weed control.

(For a complete list of terrestrial use sites, see the Directions for Use section of the attached labeling inside.)

See attached labeling inside for Complete Directions for Use in English and Spanish.

Read the entire label before using this product.

Use only according to label directions.

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

THIS IS AN END-USE PRODUCT. MONSANTO COMPANY DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Read the LIMIT OF WARRANTY AND LIABILITY statement on side panel before buying or using. If terms are not acceptable, return at once unopened.

This product is protected by U.S. Patent No.'s. 5,683,958; 5,703,015; 6,063,733; 6,121,199; 6,121,200. No license granted under any non-U.S. patent(s).

Packed for: MONSANTO COMPANY, 800 N. LINDBERGH BLVD., ST. LOUIS, MISSOURI 63167 USA

EPA Reg. No. 524-475

EPA Est. 524-IA-1

©2018 101816

Keep out of reach of children.

CAUTION See side panel for additional Precautions and First Aid.

ACTIVE INGREDIENT:
 *Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt 41.0%
 OTHER INGREDIENTS (including surfactant): 59.0%
 100.0%

*Contains 480 grams of the active ingredient glyphosate, in the form of its isopropylamine salt, per liter or 4 pounds per U.S. gallon, which is equivalent to 356 grams of the acid, glyphosate, per liter or 3 pounds per U.S. gallon (30.4% by weight).

TWO ^{2.5} GAL UNITS

Manténgase fuera del alcance de los niños.

PRECAUCION Lea las precauciones en el costado de la caja.

INGREDIENTE ACTIVO:
 *Glifosato, N-(fosfonometil) glicina, en la forma de su sal de isopropilamina 41.0%
 OTROS INGREDIENTES (incluyendo surfactante): 59.0%
 100.0%

*Contiene 480 gramos por litro o 4 libras por galón norteamericano del ingrediente activo glifosato, en la forma de su sal de isopropilamina, lo cual es equivalente a 356 gramos por litro o 3 libras por galón norteamericano (30.4% por peso) del ácido, glifosato.

DOS ^{2.5} GAL UNIDADES

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children

CAUTION

Causes eye irritation

Avoid contact with eyes or clothing

FIRST AID

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

- Have the product container or labeling with you when calling a poison control center or doctor, or going for treatment.
- You can also call (314) 694-4000, collect, day or night, for emergency medical treatment information.
- This product is identified as **Roundup PRO Herbicide, EPA Registration No. 524-475.**

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.

Physical or Chemical Hazards

Spray solutions of this product may be mixed, stored and applied using stainless steel, fiberglass, plastic or plastic-lined steel containers. DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which can form a highly combustible gas mixture. This gas mixture could flash or explode if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source and cause serious personal injury.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

SCC Barcode:
50070183888912

21136512-38

SCC Barcode:
50070183888912

MONSANTO

21136512-38

2D Code

4G/Z25.1/S/18
USA/M4389